

FINAL REPORT

HABITAT RESTORATION AT SANDY RIVER GORGE PRESERVE

1998



PROJECT DESCRIPTION

This habitat restoration project encompassed a 6.5 mile stretch of the Sandy River within the Wild and Scenic Waterway boundaries between Dodge Park and Oxbow Park. Project sites were located on The Nature Conservancy's (TNC) property, as well as, state and federal properties. The Nature Conservancy applied the grant to a contract with a ten person AmeriCorps/North West Service Academy (NWSA) team for a 16-week period to lead volunteers in the enhancement/restoration of native habitat on the Cornwell tract, maintenance of trails, expansion of community outreach and initiation of public education on appropriate land use strategies and stewardship ethics.

In early April, the NWSA team received leadership training and orientation to the history, culture and biological significance of the Sandy River Gorge. For the next 16 weeks, team members led 300 volunteers in the removal of more than 5 million Scotch Broom plants from 32 acres of land owned by TNC and Oregon Department of Fish and Wildlife (ODFW). In addition to the Scotch Broom removal, the NWSA team installed erosion barriers, improved formal trails and obliterated social trails. The NWSA team hosted volunteers from TNC, other AmeriCorps programs, volunteer youth crews, alternative education programs, school and college groups, and other community groups from as far away as Hood River.

At the beginning and the end of the project period, the NWSA team and TNC hosted picnics for people living or owning lands in the vicinity of the Sandy River Gorge. These picnics held at Oxbow Park and Dodge Park provided an opportunity for local citizens to meet the crew members, to watch demonstrations of Scotch Broom removal, to learn about other land management activities taking place in the gorge, and to discuss individual matters of interest. These gatherings had a positive impact on the local community. Attendees were eager to see work continue in future years along with annual community picnics where they could be updated on management achievements.

GOALS AND BENEFITS OF PROJECT

The primary goal and benefit was to remove a major population of Scotch Broom within the Wild and Scenic Waterway boundary. This was successfully accomplished and returned large forest meadows back to its native habitat condition. Trails that were created for public use were improved while trails impacting sensitive habitat were obliterated. Other noxious weeds were identified and removed before they could become a problem in the future. It also created an opportunity to expand our community outreach leading to involvement with the Sandy River Basin Watershed Council for future partnership endeavors.

The restoration grant also leveraged a subsequent grant through the Lower Columbia River Estuary Program to complete an initial invertebrate inventory in the Sandy River Gorge and construct informational kiosks at Dodge Park and at the entrance to TNC's Diack Tract of the Sandy River Preserve.

PROJECT DESCRIPTION

This habitat restoration project encompassed a 6.5 mile stretch of the Sandy River within the Wild and Scenic Waterway boundaries between Dodge Park and Oxbow Park. Project sites were located on The Nature Conservancy's (TNC) property, as well as, state and federal properties. The Nature Conservancy applied the grant to a contract with a ten person AmeriCorps/North West Service Academy (NWSA) team for a 16-week period to lead volunteers in the enhancement/restoration of native habitat, maintenance of trails, expansion of community outreach and initiation of public education on appropriate land use strategies and stewardship ethics.

In early April, the NWSA team received leadership training and orientation to the history, culture and biological significance of the Sandy River Gorge. For the next 16 weeks, team members led 300 volunteers in the removal of more than 5 million Scotch Broom plants from lands owned by TNC, Bureau of Land Management, METRO, Portland Water Bureau and Oregon Department of Fish and Wildlife (ODFW). On the ODFW property alone, 32 acres were cleared of mature Scotch Broom. In addition to the Scotch Broom removal, the NWSA team installed erosion barriers, improved formal trails and obliterated social trails. The NWSA team hosted volunteers from TNC, other AmeriCorps programs, volunteer youth crews, alternative education programs, school and college groups, and other community groups from as far away as Hood River.

At the beginning and the end of the project period, the NWSA team and TNC hosted picnics for people living or owning lands in the vicinity of the Sandy River Gorge. These picnics held at Oxbow Park and Dodge Park provided an opportunity for local citizens to meet the crew members, to watch demonstrations of Scotch Broom removal, to learn about other land management activities taking place in the gorge, and to discuss individual matters of interest. These gatherings had a positive impact on the local community. Attendees were eager to see work continue in future years along with annual community picnics where they could be updated on management achievements.

GOALS AND BENEFITS OF PROJECT

The primary goal and benefit was to remove the major populations of Scotch Broom within the Wild and Scenic Waterway boundary. This was successfully accomplished and returned large forest meadows back to their native habitat conditions. Trails that were created for public use were improved while trails impacting sensitive habitat were obliterated. Other noxious weeds were identified and removed before they could become a problem in the future. It also created an opportunity to expand our community outreach leading to involvement with the Sandy River Basin Watershed Council for future partnership endeavors.

The restoration grant also leveraged a subsequent grant through the Lower Columbia River Estuary Program to complete an initial invertebrate inventory in the Sandy River Gorge and construct informational kiosks at Dodge Park and at the entrance to TNC's Diack Tract of the Sandy River Preserve.

1997 METRO HABITAT RESTORATION GRANT

TASKS COMPLETED	VOLUNTEER HOURS	DATES
DIACK TRACT		
Removed English ivy from 25 trees on the Diack tract.		4/8/97
Closed Beaver Dam trail.		4/9/97
Cleared trail and rerouted section around landslide on Anne's trail.	60	4/29/97
CORNWELL TRACT		
Cleared 32 acres of Scotch broom.	2,004	4/3/97 - 7/31/97
Removed Japanese knotweed from 25' x 25' section.		5/22/97 - 7/24/97
Removed Himalayan Blackberry from riverbank		7/16/97
Set up photo point monitoring system.		6/17/97 - 7/29/97
OXBOW REGIONAL PARK		
Removed Scotch broom from Picnic Areas A and Camping Area 2.	332	4/22/97 - 5/9/97
DODGE PARK		
Removed all of Scotch broom in park.	75	6/17/97
BLM/PARTRIDGE TRACT		
Obliterate 150' of trail extending leading to ACEC	96	5/14/97 - 5/15/97
Removed 550 Scotch broom plants in clear-cut area.	96	5/21/97 - 5/22/97 & 7/17/97
Maintained trail by installing 9 drains, one 30' trench, 5 water bars, and two sets of steps. Also improved tread to encourage drainage.	132	5/8/97 & 7/17/97 - 7/18/97
TOTAL VOLUNTEER HOURS	2,795	

**HABITAT RESTORATION AT SANDY RIVER GORGE PRESERVE
STAFF, WORKKERS AND VOLUNTEERS**

Staff:

- Eddie Huckins, Portland Area Preserve Manager
- Lupine Jones, Volunteer Program Manager

Workers

- Alisa Reich, NWSA Individual Placement
- Jon Jackson, NWSA Team Leader

Team Members:

- Claudia Baskind
- Patrick Scoffield
- Frank Dotson
- Marisa Porte
- Amy Stone
- Michelle Desilets
- Jean White
- Kim McLellan
- Melissa Locke

Volunteer Organizations

- Taproot
- Multnomah Youth Cooperative
- New Trails
- C.R.U.E.
- Milwuakie Junior High
- Cascade Education Corps
- Environmental Middle School
- Mt. Hood Community College
- McGloughlin Junior High
- Gresham High School
- Oregon Episcopal School
- Reynolds High School
- ESD Re-Entry Program
- Progress Home
- National Civilian Community Corps
- I Have a Dream Foundation
- Youth Volunteer Corps
- Washington Summer Youth Program
- Portland Public School Turancy Diversion Project

HOW PROJECT RELATES TO THE GREENSPACES PROGRAM

This project clearly relates to the Greenspaces Program in that it allowed for restoration actions that would not have otherwise been completed; it increased public awareness of conditions impacting the natural landscape; it applied community-based conservation across many jurisdictions in a cooperative solution to a common problem. It is the intent of TNC and other federal/state agencies to maintain the project sites as a natural area with only passive recreational opportunities in perpetuity.

WHAT WORKED / WHAT DIDN'T / HELPFUL HINTS

Scotch Broom plants were removed manually using Weed Wrenches®, pulaskis, handsaws, and loppers. Removed plants were placed in full sun in many small to medium sized piles which allowed biomass to decompose more rapidly; piles were not evident one year later.

Japanese knotweed need to be removed in a manner that does not disturb the root structure but denies nutrients to the roots. Therefore, knotweed stems were removed from root rhizomes four times during the 16-week period. The last removal occurred during the flowering stage of the knotweed's life cycle. The timing of the last stem removal was important to deny greatest amount of nutrients to the root system. Any exposed root material was removed from the site and burned. Other vegetation was piled in the sun to decompose. This process must be continued for two to three years to kill root clump. Knotweed propagates from root segments to a greater degree than from seed. Therefore, preventing disturbance to the root system is very important in controlling its spread. A TNC element stewardship abstract is attached for information since Japanese knotweed was only recently introduced to the Sandy River Gorge as a result of the 1996 flood event.


ADVICE FOR OTHER PROJECT MANAGERS


Control of noxious weed species takes patience and perseverance. For Scotch Broom, Himalayan blackberry and Japanese knotweed, plan on 3 to 5 years of intensive labor to finally see beneficial results of your actions. Second and third years can produce overwhelming new growth.

MONITORING AND MAINTENANCE PLAN

TNC is obligated by the contract to carry-out a monitoring and evaluation program for at least four years following project completion. Volunteers will continue to follow-up with weed removal and trail maintenance activities. Working with the Sandy River Basin Watershed Council, new initiatives will be pursued to expand community involvement in noxious weed control and removal.

Sandy River Gorge





Key

= area cleared of Scotch Broom
 (32 acres including 8 acres of dense Scotch Broom)

= area not yet checked for Scotch Broom
 16 acres including 9 on South side of river

= Sandy River

Δ = trail head

• = roads/trails

• = Photo points



See photograph for visual image and photopoints

CORNWELL TRACT
Photo points

Olve hole



Sandy River

N↑

N↑

ELEMENT STEWARDSHIP ABSTRACT
for

Polygonum cuspidatum
(*Reynoutria japonica*)

Japanese knotweed
or
Mexican bamboo

To the User:

Element Stewardship Abstracts (ESAs) are prepared to provide The Nature Conservancy's Stewardship staff and other land managers with current management related information on those species and communities that are most important to protect, or most important to control. The abstracts organize and summarize data from numerous sources including literature and researchers and managers actively working with the species or community.

We hope, by providing this abstract free of charge, to encourage users to contribute their information to the abstract. This sharing of information will benefit all land managers by ensuring the availability of an abstract that contains up-to-date information on management techniques and knowledgeable contacts. Contributors of information will be acknowledged within the abstract and receive updated editions.

For ease of update and retrievability, the abstracts are stored on computer at the national office of The Nature Conservancy. Each abstract has a Nature Conservancy office or program responsible for its updating and maintenance. The address and telephone number of the office is recorded on the first page of the abstract. Anyone with comments, questions, or information on current or past monitoring, research, or management programs for the species or community described in an abstract should contact the Land Steward in the office responsible for that abstract.

This abstract is a compilation of available information and is not an endorsement of particular practices or products.

Please do not remove this cover statement from the attached abstract.

Author of this Abstract: Leslie Seiger, Department of Biology, San Diego State University, San Diego CA 92182-4614,
E-mail:lseiger@sunstroke.sdsu.edu

THE NATURE CONSERVANCY

1815 North Lynn Street, Arlington, Virginia 22209 (703) 841 5300

>== 0010 EL-CODE

POLREV

>== 0012 STEW-ABS-RESP

THE NATURE CONSERVANCY

PENNSYLVANIA CHAPTER

1211 CHESTNUT STREET, 12TH FLOOR

PHILADELPHIA, PA 19107-4122

>== 0016 PREPARER

LESLIE SEIGER

Leslie Seiger

Department of Biology

San Diego State University

San Diego CA 92182-4614

E-mail:lseiger@sunstroke.sdsu.edu

>== 0020 NAME

POLYGONUM CUSPIDATUM Sieb. & Zucc.

REYNOUTRIA JAPONICA Houtt.

[FALLOPIA BALDSCHUANICA]

Polygonum cuspidatum was classified as Reynoutria japonica by Houttuyn in 1777 and as Polygonum cuspidatum by Siebold in 1846. It was not until the early part of the 20th century that these were discovered to be the same plant (Bailey, 1990). The plant is generally referred to as Polygonum cuspidatum by Japanese and American authors and as Reynoutria japonica in the United Kingdom and Europe. Recent evidence suggests that this plant should be reclassified as Fallopiia japonica (Bailey, 1990).

>== 0050 COMMON NAME

JAPANESE KNOTWEED

MEXICAN BAMBOO

JAPANESE FLEECE FLOWER

>== 0100 DESCRIPTION

The following is based on descriptions by Fernald (1950), Muenscher (1955), Locandro (1973,1978), and Ohwi (1965).

Polygonum cuspidatum is an herbaceous perennial which forms large clumps 1-3 meters high. It is fully dioecious and can reproduce by seed and by large rhizomes which may reach a length of 5-6 meters. The stout stems are hollow and bamboo-like, extend from an erect base and are simple or little branched and glabrous with thinly membranous sheaths. Leaves are broadly ovate, truncate to cuneate at base, abruptly cuspidate, 5-15 cm long, 5-12 cm broad, with petioles 1-3 cm long. Greenish white flowers 2.5-3 mm long, densely arranged in axillary panicles; 3 styles; 8-10 stamens with longitudinally dehiscent anthers. Fruiting calyx wing-angled, 6-10 mm long. Achenes shiny black-brown, 3-4 mm long, acutely

trigonus. Male flowers have branched panicles on upright racemes with the distal end of the raceme in the highest position; individual panicles generally point up. Female flowers are drooping or decumbent with the distal end in the lowest position; individual panicles are not oriented in a particular direction. Both male and female flowers possess vestigial organs of the other sex.

Polygonum cuspidatum closely resembles *Polygonum sachalinense* (*Reynoutria sachalinensis*), an exotic species native to northern Japan and the Sakhalin Islands (Ohwi, 1965). *P. sachalinense* can be distinguished primarily by its larger size, greenish flowers and cordate leaves which gradually taper to the tip (Fernald, 1950). *Polygonum cuspidatum* is known to hybridize with *Polygonum sachalinense* and with *Fallopia baldschuanica* (Bailey, 1985, 1988). Hybrids between *Polygonum cuspidatum* and *Polygonum sachalinense* have been frequently mistaken for *Polygonum cuspidatum* in the U.K. (Bailey, 1990).

>= 1000 HABITAT

Polygonum cuspidatum is native to eastern Asia. It was introduced from Japan to the United Kingdom as an ornamental in 1825, and from there to North America in the late nineteenth century (Conolly, 1977; Patterson, 1976; Pridham and Bing, 1975). In Japan, *Polygonum cuspidatum* is widely distributed and is usually found in sunny places on hills and high mountains (Kanai, 1983; Ohwi, 1965). It is a dominant pioneer in the primary succession of volcanic slopes and is frequently a colonizer in secondary succession (Hirose, 1984).

In the U.K., *Polygonum cuspidatum* has spread extensively, occurring in virtually every 10 km square (Nature Conservancy Council, 1989). Stands range in size from individual plants to clumps more than 500 square meters (Palmer, 1990). *Polygonum cuspidatum* has also become naturalized in much of central Europe (Sukopp and Sukopp, 1988). In North America, it is widely found in the eastern U.S. and has been observed as far north as Nova Scotia and New Foundland, as far south as North Carolina, in much of the midwest and in the coastal areas of Washington and Oregon (Locandro, 1978; Patterson, 1976; Pauly, 1986). Large stands have been noted in western Pennsylvania, in particular along the banks of the Ohio and Allegheny Rivers and on the islands in these rivers (Wiegman, pers. comm.). *Polygonum cuspidatum* spreads primarily along river banks, but also grows in wetlands, waste places, along roadways, and in other disturbed areas (Beerling, 1990; Conolly, 1977; Muenscher, 1950).

Polygonum cuspidatum can thrive in a wide variety of habitats. In Japan, it can grow on volcanic soils high in sulfur and having a pH less than 4 (Conolly, 1977). In the U.S., it has been observed growing in a variety of soil types, including silt, loam, and

sand, and in soils with pH ranging from 4.5 to 7.4 (Locandro, 1973). Analyses of soils from 17 stands in Wales showed no correlation between stand size and vigor and soil characteristics. The stands studied grew in soils with a broad range of pH, organic matter and nutrients (Palmer, 1990). In Japan, *Polygonum cuspidatum* growth is slow, but steady in nutrient poor sites, and rapid in nutrient rich sites (Hirose, 1984). In areas where *Polygonum cuspidatum* has been introduced, it is found primarily in moist, unshaded habitats. Distribution maps from the U.K. show that it is generally associated with regions of high precipitation (Conolly, 1977). Locandro (1973) reports it growing on xeric as well as hydric sites in the U.S. Its distribution appears to be limited by light. It is found primarily in open sites, and its growth and abundance are depressed in shady sites (Beerling, 1991; Seiger, unpublished data).

>== 2000 BIOLOGY-ECOLOGY

Polygonum cuspidatum flowers from July to October in Japan (Ohwi, 1965) and in August and September in the U.K. and North America (Conolly, 1977; Fernald, 1950; Muenscher, 1955). It is pollinated by bees and other insects (Bailey, 1990; Locandro, 1978). Seeds appear about two weeks after flowering (personal observation) and are wind dispersed (Maruta, 1976). In Japan, reproduction in *Polygonum cuspidatum* is characterized by high seed production and low seedling survival, but plants have a very high probability of survival once established (Hirose and Tateno, 1984). However, in the U.S., U.K. and Europe, seeds do not appear to be a significant mode of reproduction. In a study of the reproductive ecology of *Polygonum cuspidatum* populations in New Jersey, Locandro (1973) found viable pollen, but noted that fertile males were rare. Seedling germination was observed in the presence of males, but no seedling survival was recorded during five years of observation. In the absence of males, females produced empty achenes. In the U.K., fruit set is very rare. This was originally attributed to the rarity of male fertile flowers (Conolly, 1977). It has since been found that there are no male fertile forms of *Polygonum cuspidatum* in England and that the pollen source is actually a hybrid between *Polygonum cuspidatum* and *Polygonum sachalinense* (Bailey, 1990). *Polygonum cuspidatum* also hybridizes with *Fallopia baldschuanica* (Bailey, 1985, 1988, 1990). In the U.S., hybrids morphologically similar to those between *Polygonum cuspidatum* and *Fallopia baldschuanica* have been grown from seeds collected in the field, but seedling establishment has not been observed in the wild (Seiger, unpublished data).

The primary mode of reproduction in the U.S., U.K. and Europe is through extensive rhizomes which can reach 15-20 meters in length (Conolly, 1977; Locandro, 1973). Dispersal can occur naturally when rhizome fragments are washed downstream by the current and deposited on banks or, as more commonly occurs, when soil is transported by humans as fill dirt (Conolly, 1977; Locandro,

1978). Rhizomes can regenerate from small fragments, and have even been observed to regenerate from internode tissue (Locandro, 1973). Rhizomes can regenerate when buried up to 1 meter deep and have been observed growing through two inches of asphalt (Locandro, 1978; Pridham and Bing, 1975). The ability of rhizomes to generate shoots was found to be affected by the source of rhizome fragments, fragment size and depth planted, the optimal depth being just below the surface (Locandro, 1973).

Polygonum cuspidatum requires high light environments and effectively competes for light in such environments by emerging early in the spring and using its extensive rhizomatous reserves to quickly attain a height of 2-3 meters. Shoots generally begin to emerge in April and growth rates exceeding 8 cm per day have been recorded (Locandro, 1973). In addition, its deep root system gives it an advantage in foraging for nutrients and water, and contributes to soil stabilization on disturbed sites (Hirose and Tateno, 1984; Nakamura, 1984). Hirose and Tateno (1984) found that organic nitrogen levels on Mt. Fuji increased following colonization by *Polygonum cuspidatum* on bare volcanic desert and concluded that *Polygonum cuspidatum* contributes to the development of the ecosystem, in part, by acting as a nutrient reservoir.

Polygonum cuspidatum is found on open sites and does not appear to be able to invade forest understory due to its high light requirements (Beerling, 1991). Studies of the very closely related *Polygonum sachalinense* indicated that *Polygonum sachalinense* plants grown in low light did not have higher photosynthetic rates than plants grown in high light, and thus would not be expected to adapt to sites with low light intensity (Patterson, Longstreth and Peet, 1977). Transplant studies of *Polygonum cuspidatum* in closed understory sites showed poor survival and growth compared to open bank sites, confirming that it was environmental factors and not limitations on dispersal which exclude *Polygonum cuspidatum* from understory sites (Seiger and Merchant, 1991). Follow-up studies in the greenhouse showed that *Polygonum cuspidatum* grown under light levels comparable to those found in the understory had significantly less rhizomatous reserves at the end of the season than did plants grown under full sunlight (Seiger, unpublished data).

>== 2500 EO-QUAL-DET

>== 3000 THREATS

Polygonum cuspidatum is widely distributed in much of the eastern U.S. In western Pennsylvania it already occupies hundreds of acres of wetlands, streambanks and hillsides and has spread along the banks of the Allegheny and Ohio Rivers and dominates the edges of many of the islands in these rivers (Wiegman, pers.comm.). It is present on at least two sites belonging to the Pennsylvania Chapter of The Nature Conservancy (Long Pond in the Poconos and Bristol Marsh, an urban preserve near Philadelphia) and has become

a problem on creeks in suburban Philadelphia (Broaddus, pers. comm.). It is also a serious problem in Rock Creek Park, a national park in Washington D.C. In the U.K., it is considered a major weed and a threat to conservation, and it is legally prohibited to introduce Japanese knotweed into the wild (Beerling, 1990; Nature Conservancy Council, 1989; Palmer, 1990). Its early emergence and great height combine to shade out other vegetation and prohibit regeneration of other species (Sukopp and Sukopp, 1988). Thus it reduces species diversity and damages wildlife habitat (Palmer, 1990; Scott and Mars, 1984; Wiegman, pers. comm.). It does not appear to be a threat in undisturbed forest and other low light areas, but it is likely that, if unchecked, it will continue to expand its range in open habitats. Once *Polygonum cuspidatum* has established it forms large, almost pure stands which are extremely persistent and difficult to eradicate.

>== 3500 LAND-PROT-SPECS

>== 4000 RECOVERY-POT

>== 5000 BIOL-MONIT-NEEDS

It is extremely difficult, if not impossible to eradicate large established stands of *Polygonum cuspidatum*. However, establishment can be prevented fairly easily by removing *Polygonum cuspidatum* before it becomes firmly entrenched. Areas known to be near established stands of *Polygonum cuspidatum*, particularly those downstream from such stands, should be monitored for the introduction of *Polygonum cuspidatum*.

>== 5200 BIOL-MONIT-PROCS

>== 5400 BIOL-MONIT-PROGS

The following individuals have direct experience monitoring *Polygonum cuspidatum*:

Leslie Seiger

Dept. Biological Sciences

The George Washington University

Washington DC 20052

(202) 994-7144

Robert Orchowski, Director, Air Quality, Environmental Affairs Unit

Duquesne Light

One Oxford Center

301 Grant Street

Pittsburgh, PA 15279

(412) 393-6099

John Palmer

Richards Moorehead & Laing Ltd

3 Clwyd Street

Ruthin

Clwyd LL15 1HF, Wales

David Beerling

School of Pure & Applied Biology
University of Wales
PO Box 915
Cardiff CF1 3TL, Wales

>== 6000 RSRCH-NEEDS

>== 6010 RSRCH-NEEDS-COMM

Current control methods (mechanical, herbicidal) require continued treatment to prevent reestablishment of Polygonum cuspidatum. It may be feasible to reintroduce competitors as an alternative to continued treatment. There is a need for more research on whether native species might serve effectively as competitors and methods of reintroduction. Only very preliminary work has been done towards developing a biological control for Polygonum cuspidatum and much research remains to be done (see below).

>== 6400 RSRCH-PROGS

>== 6410 RSRCH-PROGS-COMM

The International Institute for Biological Control in conjunction with the National Agricultural Research center in Japan has produced a partial list of insect herbivores in Japan which are associated with Polygonum cuspidatum. A number of pathogens have been collected from Japan by CAB International and are held at the International Mycological Institute (Fowler and Schroeder, 1990). Plans are under way to begin a study of Polygonum cuspidatum in its native habitat (Fowler, pers. comm.). Contact:

Dr. Simon V. Fowler
International Institute of Biological Control
Silwood Park
Buckhurst Road
Ascot, Berks, SL5 7TA, UK

Recent studies in Rock Creek park, Washington, D.C. indicate potential for control by mechanical means combined with revegetation (Seiger, unpublished data). Field tests will be conducted in 1992. Contact:

Leslie Seiger
Dept. Biological Sciences
The George Washington University
Washington, DC 20052
(202)994-7144

>== 7000 MGMT-NEEDS

>== 7010 MGMT-NEEDS-COMM

In areas where Polygonum cuspidatum has not yet become established, the focus of management should be to prevent establishment by monitoring areas for introductions of Polygonum cuspidatum and eradicating newly established stands before they can become established.

>== 7400 MGMT-PROCS

Manual control consists of digging out the rhizomes or cutting the stalks. Digging is extremely labor intensive and tends to spread the rhizome fragments and promote disturbance and is not recommended (Palmer, 1990). Cutting, on the other hand, may be quite effective in eliminating *Polygonum cuspidatum*. It has been observed that *Polygonum cuspidatum* does not establish where grazing pressure is high (Beerling, 1990; Palmer, 1990). In a review of control methods, Palmer (1990) noted that eradication is not complete with cutting alone, but has been nearly achieved in some cases and should be feasible with persistence. A number of authors claim that cutting is ineffective (Pauly, 1986; Pridham and Bing, 1975; Orchowski, 1991). These conclusions are based on the observation that shoots are regenerated following cutting. However, a greenhouse study found that cutting stems results in a significant reduction of rhizomatous reserves. The same study also found that cutting was equally effective at any time during the growing season prior to the beginning of senescence (Seiger and Merchant, 1990). A study of the effects of repeated cutting showed that at least three cuts are needed in a growing season to offset rhizome production (Seiger, unpublished data). Manual control can be labour intensive, but where populations are small and isolated, may be the best option. No research has been done to test the effectiveness of burning. It may act similarly to cutting by removing above ground material.

Shading, particularly in conjunction with cutting, may be another useful means of control on smaller stands. Studies showing that *P. cuspidatum* requires high light environments suggest that covering stands with black plastic or shade cloth may reduce growth. Pridham and Bing (1975) state that applying several layers of black polyethylene film covered by asphalt, blocks or stones to a leveled soil surface may provide some control. However, they also note that *P. cuspidatum* is able to emerge through asphalt. If shade cloth (or plastic) is to be applied without cutting, then, to prevent *P. cuspidatum* from emerging through the covering, shade cloth should be placed over shoots after the plants have reached their full height or placed well above newly emerging shoots, or raised as plants grow.

A number of biocidal chemicals have been found to be effective against *Polygonum cuspidatum*. Most of these are undesirable for use in conservation areas because they are nonselective, may be persistent in the soil and/or are not safe for use near water. One frequently used way to minimize the effects of non-selective herbicides on non-target species is to paint herbicides directly onto the target plants (Broadus, pers. comm.). In the case of *P. cuspidatum*, this would probably require prior cutting for easier access if herbicides are to be applied after the plants have reached their full height. Herbicides appear to be more effective

when combined with cutting (Scott and Mars, 1984; Orchowski, 1991).

Glyphosate [N-(phosphonomethyl)glycine] has been found to be very effective against *Polygonum cuspidatum* (Ahrens, 1975; Beerling, 1990; Pauly, 1986). Glyphosate is a nonselective systemic herbicide with a short residual life (Ahrens, 1975; Lynn, Rogers and Graham, 1979). Application is more effective in the fall when leaves are translocating to rhizomes (Lynn, et al, 1979). The British Nature Conservancy Council (1989) recommends applying 2.0 kg/ha in August with a prior cut in late spring or early summer. Glyphosate is available from Monsanto under the trade names Roundup™ and Rodeo™. Only Rodeo has been approved for use near water (Bender, 1988). Glyphosate has been used with limited success on some nature reserves in the U.K. (Palmer, 1990). Repeated applications over several years may be necessary (Beerling, 1990; Palmer, 1990; Pauly, 1986).

The Nature Conservancy Council (1989) also recommends picloram to be applied at a rate of 2.6 kg/ha in the spring. Picloram is a selective herbicide which is persistent in the soil. It must not be used near water, thus excluding its use in many of the areas where *Polygonum cuspidatum* is a problem (Gritten, 1990; Scott and Mars, 1984). Dicamba (3,6-dichloro-o-anisic acid) has also been found to be effective against *Polygonum cuspidatum*, but is persistent in the soil and nonselective (Pridham and Bing, 1975). A number of other herbicides have been tested against *Polygonum cuspidatum*, both alone and in combination with other herbicides (Orchowski, 1991; Scott and Mars, 1984). Herbicide may have to be used on stands that have been allowed to attain a large size. However, their use is not recommended in nature reserves because of their undesirable effects on other biota and the need for repeated applications to maintain control of *Polygonum cuspidatum*.

Regardless of whether control is manual or chemical, as long as some rhizomes remain in the soil *Polygonum cuspidatum* returns once management is relaxed (Beerling, 1990; Nature Conservancy Council, 1989; Palmer, 1990). It has been suggested that the reintroduction of effective competition might be possible (Eaton, 1986).

Research has only recently begun on biological control. The herbivores and pathogens of *Polygonum cuspidatum* in Wales have been examined for their potential as control agents (Fowler and Schroeder, 1990). A program is underway at the International Institute of Biological Control to identify biological control agents (Fowler, pers. comm.). The genetic uniformity of *Polygonum cuspidatum* makes it a good candidate for biological control (Bailey, 1990). Biological control may be necessary where *Polygonum cuspidatum* has taken over vast areas as it has done in the U.K., but it may be years before a successful control agent can be found. >== 7700 MGMT-PROGS

>== 7710 MGMT-PROGS-COMM

The following individuals are familiar with *Polygonum cuspidatum* and its control in natural areas:

Leslie Seiger
The Dept. of Biological Sciences
The George Washington University
Washington, D.C. 20052
(202)994-7144

John Palmer
Richards Moorehead & Laing Ltd
3 Clwyd Street
Ruthin
Clwyd LL15 1HF, Wales

>== 8000 SUM-STEW-NEEDS

Polygonum cuspidatum occurs in much of the temperate U.S. Though not yet a major weed in the U.S., it is spreading, particularly in the eastern states. Dispersal is limited to areas where rhizome fragments from existing stands are washed downstream or soil containing rhizomes is transported by humans. Once established, it forms large, monospecific stands which displace virtually all other vegetation. Establishment can be prevented by monitoring for the introduction of *Polygonum cuspidatum* and manually removing the entire plant. Small stands may be controlled by repeated cutting, which may need to be supplemented by revegetation once growth of *Polygonum cuspidatum* has been reduced. At present, the only method to control large stands is with repeated application of herbicides. Complete eradication may not be possible.

>== BIBLIOGRAPHY

Ahrens, J.F. (1975). Preliminary results with glyphosate for control of *Polygonum cuspidatum*. Proc. Northeast. Weed Science Soc. 29:326.

Bailey, J.P. (1985). Chromosome numbers of some alien *Reynoutria* species in the British Isles. *Watsonia*, 15:269-277.

Bailey, J.P. (1988). Putative *Reynoutria japonica* Houtt. x *Fallopia baldschuanica* (Regel) Holub hybrids discovered in Britain. *Watsonia*, 17:163-181.

Bailey, J.P. (1990). Breeding behaviour and seed production in Alien Giant Knotweed in the British Isles. In: The biology and control of invasive plants. Conference organized by the Industrial Ecology Group of the British Ecological Society at the University of Wales College of Cardiff, September 20-21, 1990.

Beerling, D.J. (1990). The use of non-persistent herbicides to control riparian stands of Japanese Knotweed (*Reynoutria Japonica*

Houtt). In: The biology and control of invasive plants. Conference organized by the Industrial Ecology Group of the British Ecological Society at the University of Wales College of Cardiff, September 20-21, 1990.

Beerling, D.J. (1991). The effect of riparian land use on occurrence and abundance of Japanese knotweed *Reynoutria japonica* on selected rivers in South Wales. *Biol. Cons.* 55:329-337.

Bender, J. (1988). Element Stewardship Abstract for *Lythrum salicaria*. The Nature Conservancy.

Broadus, L. (1992). The Nature Conservancy, Pennsylvania Chapter. Memorandum to Leslie Seiger, March 6, 1992.

Conolly, A.P. (1977). The distribution and history in the British Isles of some alien species of *Polygonum* and *Reynoutria*. *Watsonia* 11:291-311.

Eaton, J.W. (1986). Waterplant ecology in landscape design. In *Ecology and design in landscape*. The 24th symposium of the British Ecological Society, Manchester 1983. A.D. Bradshaw, D.A. Goode, E.H.P. Thorp (Eds.). Blackwell Scientific Publications. Oxford.

Fernald, M.L. (1950). *Gray's Manual of Botany*. 8th Ed. American Book Company, N.Y.

Fowler, S.V. (1991). International Institute of Biological Control. Silwood Park, Buckhurst Road, Ascot, Berks, SL5 7TA, UK. Letter to Leslie Seiger, January 8, 1991.

Fowler, S.V. and D. Schroeder (1990). Biological control of invasive plants in the UK - prospects and possibilities. In: The biology and control of invasive plants. Conference organized by the Industrial Ecology Group of the British Ecological Society at the University of Wales College of Cardiff, September 20-21, 1990.

Gritten, R. (1990). The control of invasive plants in the Snowdonia National Park. In: The biology and control of invasive plants. Conference organized by the Industrial Ecology Group of the British Ecological Society at the University of Wales College of Cardiff, September 20-21, 1990.

Hirose, T. (1984). Nitrogen use efficiency in growth of *Polygonum cuspidatum* Sieb. et Zucc. *Annals of Botany*, 54:695-704.

Hirose, T., and M. Tateno (1984). Soil nitrogen patterns induced by colonization of *Polygonum cuspidatum* on Mt. Fuji. *Oecologia* 61:218-223.

Kanai, H. (1983). Study on the distribution patterns of Japanese plants (5): Accumulation of phytogeographical data of popular plants, points and measures. *Journ. Jap. Bot.*, 59: 257-269.

Locandro, R.R. (1973). Reproduction ecology of *Polygonum cuspidatum*. Ph.D. thesis. Rutgers University.

Locandro, R.R. (1978). Weed watch: Japanese bamboo - 1978. *Weeds Today* 9:21-22.

Lynn, L.B., R.A. Rogers, and J.C. Graham (1979). Response of woody species to glyphosate in northeastern United States. *Proc. Northeast Weed Sci. Soc.* 33:336-342.

Maruta, E. (1976). Seedling establishment of *Polygonum cuspidatum* on Mt. Fuji. *Jap. J. Ecol.* 26:101-105.

Maruta, E. (1983). Growth and survival of current-year seedlings of *Polygonum cuspidatum* at the upper distributional limit on Mt. Fuji. *Oecologia* 60:316-320.

Muenschler, W.C. (1955). *Weeds*. Second edition. McMillan Press, New York.

Nakamura, T. (1984). Vegetational recovery of landslide scars in the upper reaches of the Oi River, Central Japan. *J. Jap. For. Soc.* 66(8):328-332.

Nature Conservancy Council (1989). Japanese knotweed. In *Urban Wildlife News*, 6(2):3.

Ohwi, J. (1965). *Flora of Japan*. Smithsonian Institute, Washington, DC.

Orchowski, R.W. (1991). Knotweed control pilot program Brunot Island Wildlife Habitat Enhancement Project. Presented at the 3rd Annual Wildlife Enhancement Council Symposium, Washington, DC, November 19, 1991.

Palmer, J.P. (1990). Japanese knotweed (*Reynoutria japonica*) in Wales. In: *The biology and control of invasive plants*. Conference organized by the Industrial Ecology Group of the British Ecological Society at the University of Wales College of Cardiff, September 20-21, 1990.

Patterson, D.T. (1976). The history and distribution of five exotic weeds in North Carolina. *Castanea* 41:177-180.

Patterson, D.T., D.J. Longstreth, and M.M. Peet (1977). Photosynthetic adaptation to light intensity in Sakhalin knotweed (*Polygonum sachalinense*). *Weed Science* 25:319-323.

Pridham, A.M.S. and A. Bing (1975). Japanese-bamboo plants. *Gard.* 31:56-57.

Pauly, W.R. (1986). Summary of Mexican bamboo control methods (Wisconsin). *Restoration and Management Notes*, 4(1):37-38.

Scott, R., and R.H. Mars (1984). Impact of Japanese knotweed and methods of control. *Aspects of applied biology* 5, Weed control and vegetation management in forestry and amenity areas pp. 291-296.

Seiger, L.A. and H.C. Merchant (1990). The ecology and control of *Polygonum cuspidatum*. *Bull. Ecol. Soc. Am.* 71(2 suppl.):322.

Seiger, L.A. and H.C. Merchant (1991). Effects of site on survivorship and size of *Polygonum cuspidatum*. *Bull. Ecol. Soc. Am.* 72(2 suppl.):247.

Sukopp, H. and U. Sukopp (1988). *Reynoutria japonica* Houtt. in Japan und in Europa. *Veroff. Geobot. Inst. ETH, Stiftung Rubel, Zurich* 98:354-372.

Wiegman, P. (1992). Director of natural science and stewardship, Western Pennsylvania Conservancy, Pittsburgh, PA. Telephone conversation with Leslie Seiger, February 25, 1992.

>== 9999 END



PRINT 4 ROLL 00320162 19JUN97

PRINT 6 ROLL 00320162 19JUN97

SEATTLE FILMWORKS CALL 1-800-FILMWORKS

PRINT 16 ROLL 61839577 08AUG97

PRINT 21 ROLL 00320162 19JUN97



PRINT 9 ROLL 00312912 25JUL97

PRINT 7 ROLL 00312912 25JUL97

PRINT 11 ROLL 00320162 19JUN97



PRINT 3 ROLL 00320162 19JUN97

PRINT 7 ROLL 00320162 19JUN97

SEATTLE FILMWORKS CALL 1-800-FILMWORKS

PRINT 15 ROLL 61839577 08AUG97

PRINT 20 ROLL 00320162 19JUN97



SEATTLE FILMWORKS CALL 1-800-FILMWORKS

PRINT 21 ROLL 61722876 27JUN97

SEATTLE FILMWORKS CALL 1-800-FILMWORKS

PRINT 19 ROLL 61722876 27JUN97

PRINT 12 ROLL 00320162 19JUN97